arXiv:1503.08794v1 [gr-qc] 30 Mar 2015

From Causal Dynamical Triangulations To Astronomical Observations[a]

Jakub Mielczarek[1,2,3]
[1]Laboratoire de Physique Subatomique et de Cosmologie, UJF, INPG,
CNRS, IN2P3 53, avenue des Martyrs, 38026 Grenoble cedex, France
[2]Institute of Physics, Jagiellonian University, Łojasiewicza 11, 30-348 Cracow, Poland
[3]Department of Fundamental Research, National Centre for Nuclear Research,
Hoża 69, 00-681 Warsaw, Poland[†]

This essay discusses phenomenological aspects of the diffusion time dependence of the spectral dimension predicted by the Causal Dynamical Triangulations (CDT) approach to quantum gravity. The deformed form of the dispersion relation for the fields defined on the CDT space-time is reconstructed. Using the *Fermi* satellite observations of the GRB 090510 source we find that the energy scale of the dimensional reduction is $E_* > 6.7 \cdot 10^{10}$ GeV at (95 % CL).

By applying the deformed dispersion relation to the cosmological perturbations it is shown that, for a scenario when the primordial perturbations are formed in the UV region, the scalar power spectrum $\mathcal{P}_S \propto k^{n_S-1}$ where $n_S - 1 \approx \frac{3r(d_{\rm UV}-2)}{r+48(d_{\rm UV}-3)}$. Here, $d_{\rm UV} \approx 2$ is obtained from the CDT value of the spectral dimension in the UV limit and r is the tensor-to-scalar ratio. We find that, the predicted deviation from the scale-invariance ($n_S = 1$) is in contradiction with the up to date *Planck* and *BICEP2* results.

---

[a] Essay written for the Gravity Research Foundation 2015 Awards for Essays on Gravitation.

Submitted on 30 February 2015.

[†] jakub.mielczarek@uj.edu.pl

Typeset by REVTEX

I. INTRODUCTION

The littleness of the Planck length is indescribable. The immense size the observable Universe is paralyzing. However, the geometric mean of the two characteristic values results in a tangible quantity. Explicitly, $\sqrt{1 l_{\rm Pl} \cdot 1{\rm Gpc}} \approx 1{\rm mm}$, where the Planck length $l_{\rm Pl} \approx 1.62 \cdot 10^{-35}$ m and the gigaparsec $1{\rm Gpc} \approx 3.09 \cdot 10^{28}$ m.

As we will show in the example discussed in the essay, better understanding of both the super-short and super-large scales can be gained thanks to this property.

So, how does it work? There are two main possibilities. The first one is a result of accumulation of the tiny Planck scale deviations acting on particles propagating across cosmological distances. The second possibility exploits the evolution of the Universe. The Universe is as big as it is because it is expanding. As a result, the Planck scale in the early universe, multiplied by the growth factor becomes macroscopic in the mature universe. In both cases, the multiplication law brings the Planck scale closer to the scales of our perception.

We do not know yet what the Planck scale physics is. However, by taking the use of the compensation of scales discussed above, one may already try to falsify the theoretical models of quantum gravitational phenomena[1], which we have.

The theory of Planck scale physics which we are going to examine here is Causal Dynamical Triangulations (CDT) [1, 2]. The theory is a non-perturbative approach based on the path integral formulation of quantum mechanics. The crucial ingredient of the theory is a causality condition imposed at the level of Wick-rotated time coordinate. In similarity to statistical physics, the different space-time configurations are evaluated using Monte Carlo computer simulations.

Various promising results arose from numerical studies of the quantum gravitational system with the positive cosmological constant. One of the most profound is the emergence of four dimensional de Sitter space-time [3, 4] in the so-called phase C[2]. However, as reflected by features of the process of diffusion taking place at the simplicial manifold of CDT, the reconstructed de Sitter space-time is not fully classical.

---

[1] Which are expected to occur at the Planck scale.

[2] Existence of the three different phases of CDT (A, B and C) has been firmly confirmed so far [5].

In particular, it has been shown that the spectral dimension[3] characterizing the diffusion process can be parametrized in the following way [6]:

$$d_S(\sigma) = a - \frac{b}{c+\sigma}, \tag{1}$$

where σ is the diffusion time. At the representative point in phase C ($\kappa_0 = 2.2$, $\Delta = 0.6$) the values $a = 4.02$, $b = 119$ and $c = 54$ [6] and $a = 4.06$, $b = 135$ and $c = 67$ [7] have been obtained from numerical simulations. In both cases, four dimensional space-time ($d_S \approx 4$) is correctly recovered in the IR limit ($\sigma \to \infty$). However, in the UV limit ($\sigma \to 0$) dimensional reduction to $d_S \approx 2$ is predicted. This effect is not reserved to CDT only and has been noticed in other approaches to quantum gravity as well [8].

The expression (1) is a starting point for further considerations. After fixing the IR value d_S to be precisely equal to the topological space-time dimension $d = 4$ and parametrizing the departure from $d_S = 2$ in UV by ϵ we obtain

$$d_S(\sigma) = 4 - \frac{2-\epsilon}{1+\sigma/c}, \tag{2}$$

such that $d_{\text{UV}} := d_S(\sigma \to 0) = 2 + \epsilon$. The parameter c can be related to the energy scale of the dimensional reduction: $E_* := \frac{1}{\sqrt{c}}$.

II. FROM THE SPECTRAL DIMENSION TO THE DEFORMED DISPERSION RELATION

The spectral dimension discussed in the previous section is formally defined as follows

$$d_S(\sigma) \equiv -2\frac{\partial \log P(\sigma)}{\partial \log \sigma}, \tag{3}$$

where

$$P(\sigma) = \int d\mu e^{\sigma \triangle_p} \tag{4}$$

---

[3] The precise definition will be given in the subsequent section.

is the average return probability of the random walk (diffusion) process. Here, $d\mu$ is the invariant[4] measure in the momentum space, which we assume to be the classical one[5] $d\mu = \frac{dE d^3\vec{p}}{(2\pi)^4}$. $\triangle_p$ is the Laplace operator in momentum space. In the case of the Euclidean 4-dimensional space $\triangle_p = -E^2 - \vec{p}^2$. This expression might be viewed as the Wick rotated $(E \to iE)$ version of the analogous formula in the Minkowski space $\triangle_p = E^2 - \vec{p}^2$, which defines dispersion relation for the free fields living on the space-time ($\triangle_p = 0 \Rightarrow E^2 = \vec{p}^2$). In this case, the equations (3) and (4) predict $d_S = d$, where $d = 4$ is the topological dimension.

CDT predicts that the spectral dimension varies as a function of the diffusion time σ. The corresponding moment Laplace operator has to be, therefore, deformed with respect to the classical case. Here, we parametrize this departure in the following manner

$$\triangle_p = -E^2 - \Omega(p)^2, \tag{5}$$

where $\Omega(p)$ is some unknown function of $p := \sqrt{\vec{p} \cdot \vec{p}}$. The task would be now to recover the form of the function $\Omega(p)$ for the CDT spectral dimension (2). In this case, the expression for the return probability is

$$P(\sigma) = \frac{1}{\sqrt{4\pi\sigma}} \frac{4\pi}{(2\pi)^2} \int_0^\infty dp\, p^2 e^{-\sigma\Omega^2(p)}, \tag{6}$$

where integrations over E and the angular part of $d^3\vec{p}$ have been performed. As it has been shown in Ref. [10], the relation (6) can be converted into the form of the inverse Laplace transform

$$p^3 = \frac{1}{2\pi i} \lim_{T\to\infty} \int_{\gamma - iT}^{\gamma + iT} \left(12\pi^{5/2} \frac{P(\sigma)}{\sqrt{\sigma}} \right) e^{\sigma\Omega^2(p)} d\sigma, \tag{7}$$

the $\gamma \in \mathbb{R}$ has to satisfy the condition $\forall_{\sigma_*} \gamma > \Re(\sigma_*)$, where σ_* denotes singularities of $P(\sigma)/\sqrt{\sigma}$. The $P(\sigma)$ function is recovered by integrating definition (3) with use of parametrization (2). We find that

$$P(\sigma) = \frac{1}{16\pi^2 \sigma^{1+\epsilon/2} (\sigma + c)^{1-\epsilon/2}}, \tag{8}$$

---

[4] With respect to the isometries of a given space-time.

[5] With this assumption we enforce that the momentum space is flat as in the classical case. However, as it is hypothesized *e.g.* in the context of Relative Locality [9], non-vanishing curvature of the momentum space may be characteristic for the the quantum gravitational phenomena. This issue in the context of CDT will be studied elsewhere.

which, when applied to (7), leads to

$$p^3 = \Omega^3 {}_1F_1\left(1-\epsilon/2, 5/2; -c\Omega^2\right), \tag{9}$$

where ${}_1F_1(a,b;z)$ is the confluent hypergeometric function. Equation (9), is an entangled form of the deformed dispersion relation $E = \Omega(p)$. The function $\Omega(p)$ for $\epsilon = 0$ ($d_{\mathrm{UV}} = 2$) is shown in Fig. 1.



FIG. 1. A plot of the dispersion relation predicted by dimensional reduction in Causal Dynamical Triangulations. The IR (10) and UV (11) approximations are represented by the dashed (red) and the dotted (blue) lines respectively.

In the IR limit ($p \to 0$) the dispersion relation is approximated by

$$\Omega_{\mathrm{IR}}(p) \approx p + \frac{E_*}{15}(2-\epsilon)\left(\frac{p}{E_*}\right)^3, \tag{10}$$

so the classical limit is correctly recovered. However, when passing to the high energy range (the UV limit) the dispersion relation undergoes deflection to the form

$$\Omega_{\mathrm{UV}}(p) \approx \frac{2}{3}E_*\left(\frac{p}{E_*}\right)^{3-3\epsilon} \propto p^{3-3\epsilon}. \tag{11}$$

Both of the obtained limiting behaviors of the deformed dispersion relation will now be used to construct the relation of CDT to astrophysical and cosmological observations.

III. ASTROPHYSICS

The dispersion relation derived in the previous section can now be used to find the group velocity of photons. For the particles having energy $E \ll E_*$ the IR limit approximation (10) may be applied, leading to the quadratic deviation

$$v_{\text{gr}} := \frac{\partial E}{\partial p} = \frac{\partial \Omega(p)}{\partial p} \approx 1 + \frac{3}{15}(2-\epsilon)\left(\frac{E}{E_*}\right)^2 . \tag{12}$$

This expression predicts that the group velocity is increasing with the energy (superluminal behavior). Therefore, when a bunch of photons of different energies are emitted simultaneously the ones with the highest energy will arrive first to the observer. This effect can be quantified by the arrival time difference between E and $E \to 0$ energy photons. With use of formula (12), the corresponding expression is $\tau \simeq -L\frac{3}{15}(2-\epsilon)\left(\frac{E}{E_*}\right)^2$, where L is the distance to a source. Although a huge value of the energy scale E_* is expected[6], the multiplication by a sufficiently large distance L may bring the value of τ close to the observational window.

The effect can be constrained with use of the signals from the high-energy astrophysical sources such as gamma ray bursts (GRB) [11]. In particular, the GRB 090510 remote by $L \approx 5,8$ Mpc may be used. Using the constraint from the *Fermi*-Large Area Telescope [12] one can find that the energy scale of the dimensional reduction is $E_* > 6.7 \cdot 10^{10}$ GeV at (95 $\%$ CL). This constraint is definitely much stronger than any obtained from the accelerator physics experiments (which are reaching energies of the order of 10^4 GeV). However, because of the quadratic (in energy) from of the effect, the constraint is distant from the Planck scale[7].

IV. COSMOLOGY

Because of the quadratic nature of the IR variation of the group velocity, perhaps the more promising is an application of dimensional reduction to cosmology. Here, instead of the IR, the UV part of the dispersion relation will play a crucial role. The potential relevance of the UV scaling of the modified dispersion relation in cosmology has been suggested in Ref. [13]. It has been shown that dimensional reduction to $d_{\rm UV} = 2$ leads to the scale invariant vacuum fluctuations of the primordial perturbations in the UV domain. What has not been shown is how the shape of the spectrum changes when we try to relate it with the one that is observationally relevant. Here, we show that a deviation from $d_{\rm UV} = 2$ leads to a tilt in the power spectrum. This prediction is confronted with the observations of the cosmic microwave background radiation (CMB).

---

[6] Presumably of the order of the Planck energy $E_{\text{Pl}} \approx 1.22 \cdot 10^{19}$GeV.

[7] If the effect had been linear in energy, the observational constraint would be much stronger.

The resulting CDT dispersion relation $E = \Omega(p)$ can be incorporated into the equation of modes for the cosmological perturbations in the following way

$$\frac{d^2}{d\tau^2} f_k + \left(a^2\Omega^2(k/a) - \frac{z''}{z} \right) f_k = 0. \tag{13}$$

Here, $k = pa$ (a is the scale factor) and the classical limit $a^2\Omega^2(k/a) \to a^2(k/a)^2 = k^2$ is recovered correctly for $p \to 0$. The value of z depends on which kind of perturbations is considered. For the gravitational waves $z = z_T := a$ and for the scalar perturbations $z = z_S := a\frac{\dot{\phi}}{H}$. In the sub-Hubble limit $\left(a^2\Omega^2(k/a) \gg \frac{z''}{z} \right)$, the vacuum normalization of the f_k mode functions leads to

$$|f_k|^2 = \frac{1}{2a\Omega(k/a)}. \tag{14}$$

Applying this to the definition of the scalar power spectrum we find that for $k/a \gg E_*$

$$\mathcal{P}_S(k) := \frac{k^3}{2\pi^2} \frac{|f_k|^2}{z_S^2} = \frac{1}{\pi(1+w)} \left(\frac{E_*}{E_{\rm Pl}} \right)^2 \left(\frac{1}{E_*} \frac{k}{a} \right)^{3\epsilon} \propto k^{n_S - 1}, \tag{15}$$

where $n_S = 1 + 3\epsilon = 1 + 3(d_{\rm UV} - 2)$. Furthermore, we assume that the universe is filled with barotropic matter: $P = w\rho$, with $w = const$.

Indeed, for $d_{\rm UV} = 2$ the power spectrum (15) becomes scale invariant $\mathcal{P}_S(k) = \frac{1}{\pi(1+w)} \left(\frac{E_*}{E_{\rm Pl}} \right)^2 = const$. While it is tempting to relate the obtained spectrum with that inferred from the CMB nearly scale-invariant spectrum of the primordial perturbations, it is not allowed to do so directly. The spectrum (15) corresponds to the UV regime ($k/a \gg E_*$) which has to be suitably converted into the super-Hubble spectrum probed observationally. In general, this would require evolving the modes starting from the UV domain until crossing the Hubble horizon. This will unavoidably affect the shape of the power spectrum. A scenario in which knowledge of the intermediate evolution is not required corresponds to the case when the Hubble horizon is located in the UV domain. In that case *freezing* of the modes occurs already at the level of the UV spectrum (15).

In this particular situation the resulting tensor-to-scalar ratio

$$r := \frac{\mathcal{P}_T(k_0)}{\mathcal{P}_S(k_0)} = \frac{64\pi G \frac{k^3}{2\pi^2}\frac{|f_k|^2}{z_T^2}}{\frac{k^3}{2\pi^2}\frac{|f_k|^2}{z_S^2}} = 64\pi G \left(\frac{z_S}{z_T}\right)^2 = 24(1+w), \tag{16}$$

can be directly related to the CMB data[8]. The best current observational bound at the pivot scale $k_0 = 0.05\ \mathrm{Mpc}^{-1}$ is $r < 0.12$ (BICEP2/*Keck Array* and *Planck*) [14], which implicates that $w < -0.995$ when the perturbations were formed. The universe had to, therefore, be very close to the de Sitter phase when the perturbations were formed. Cosmic inflation is still needed!

The value of the power spectrum accessible observationally is at the horizon-crossing. In our case, the horizon-crossing condition $a^2\Omega^2(k/a) = \frac{z''}{z}$ translates into $k \simeq aE_* \left(\frac{3}{2}\frac{H}{E_*}\right)^{\frac{1}{3(1-\epsilon)}} := k_H$[9] for $H > E_*$. Using this, the spectral index of scalar perturbations at the horizon-crossing is

$$n_S - 1 \equiv \frac{d\ln \mathcal{P}_S(k = k_H)}{d\ln k} \approx \frac{3\epsilon r}{r + 48(\epsilon - 1)}. \tag{17}$$

The scale invariance is still recovered for $\epsilon = 0$. However, by comparing the prediction with the *Planck* satellite measurement of the spectral index ($n_S = 0.9616 \pm 0.0094$) [15] and the constraint on the tensor-to-scalar ratio [14] we find that

$$\epsilon > 0.835. \tag{18}$$

The value is in overt contradiction with the CDT-motivated assumption that $\epsilon \approx 0$. The scenario considered can be, therefore, ruled out based on the CMB data. Other scenarios, such as those with $H < E_*$ at the horizon-crossing have to be studied separately, taking into account the evolution of modes in the intermediate energy range.

---

[8] This is possible because of growth of a physical distance in the expanding universe ($\lambda \propto a$).

[9] Because of the deformed form of the dispersion relation, the condition differs from the classical expression $k \simeq aH$.

V. CONCLUSIONS

The results presented in this essay neither prove nor disprove Causal Dynamical Triangulations. What has been essentially shown is that phenomenology of CDT can be extracted from numerical studies of the spectral dimension. The predicted energy-dependence of the speed of light allowed us to rule out a possibility of low-energy dimensional reduction. Furthermore, by incorporating the predictions regarding the UV behavior of the spectral dimension to primordial cosmology, we found that the CDT-motivated cosmological scenarios may be observationally falsifiable. Moreover, we explicitly ruled out one of them with use of the observations of the CMB.

As Thomas A. Edison once said 'I have not failed. I've just found 10000 ways that won't work.' Quoting the sentence, we are convinced that unabated stubbornness in eliminating Planck scale scenarios will eventually lead to a discovery comparable only with Edison's one.

ACKNOWLEDGEMENTS

JM is supported by the Grant DEC-2014/13/D/ST2/01895 of the National Centre of Science. The author would like to thank to Daniel Coumbe for his valuable comments.

---

[1] J. Ambjorn, J. Jurkiewicz and R. Loll, Phys. Rev. Lett. **85** (2000) 924.

[2] J. Ambjorn, A. Goerlich, J. Jurkiewicz and R. Loll, Phys. Rept. **519** (2012) 127.

[3] J. Ambjorn, J. Jurkiewicz and R. Loll, Phys. Rev. Lett. **93** (2004) 131301.

[4] J. Ambjorn, A. Görlich, J. Jurkiewicz and R. Loll, Phys. Rev. Lett. **100** (2008) 091304.

[5] J. Ambjorn, S. Jordan, J. Jurkiewicz and R. Loll, Phys. Rev. D **85** (2012) 124044.

[6] J. Ambjorn, J. Jurkiewicz and R. Loll, Phys. Rev. Lett. **95** (2005) 171301.

[7] D. N. Coumbe and J. Jurkiewicz, arXiv:1411.7712 [hep-th].

[8] S. Carlip, AIP Conf. Proc. **1196** (2009) 72.

[9] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman and L. Smolin, Phys. Rev. D **84** (2011) 084010.

[10] T. P. Sotiriou, M. Visser and S. Weinfurtner, Phys. Rev. D **84** (2011) 104018.

[11] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos, S. Sarkar and , Nature **393** (1998) 763.

[12] V. Vasileiou, A. Jacholkowska, F. Piron, J. Bolmont, C. Couturier, J. Granot, F. W. Stecker and J. Cohen-Tanugi *et al.*, Phys. Rev. D **87** (2013) 12, 122001.

[13] G. Amelino-Camelia, M. Arzano, G. Gubitosi and J. Magueijo, Phys. Rev. D **87** (2013) 12, 123532.

[14] P. A. R. Ade *et al.* [BICEP2 and Planck Collaborations], [arXiv:1502.00612 [astro-ph.CO]].

[15] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571** (2014) A16.